



I'm not robot



**I am not robot!**

$\beta$ , Epoxy, 2a, 4, 7 $\beta$ , 10 $\beta$ , 13a hexahydroxytaxenone 4, diacetate benzoate ester with (2R,3S) N-benzoylphenylisoserine Paclitaxel has the following structural formula: Paclitaxel is a white to off-white crystalline powder with the empirical formula C<sub>47</sub>H<sub>51</sub>NO and a molecular weight of 660.54. It is highly lipophilic, insoluble in water, and melts at approximately 170 °C to 180 °C. Abbreviated Package Insert. The most common adverse reactions resulting in permanent discontinuation of ABRAXANE are neutropenia (3%), thrombocytopenia (3%), and peripheral neuropathy (1%). The most common side effects of ABRAXANE in people with non-small cell lung cancer include: low red blood cell count (anemia), decreased white blood cell count. See Product Monograph for complete product information. INDICATIONS. ABRAXANE is a prescription medicine used to treat: advanced breast cancer in people who have already received certain other medicines for their cancer. Appropriate management of therapy and ABRAXANE for Injectable Suspension (paclitaxel protein-bound particles for injectable suspension) is an antimicrotubule agent that promotes the assembly of microtubules. ABRAXANE monotherapy is indicated for the treatment of metastatic breast cancer in adult patients who have failed first-line treatment for metastatic disease and for whom. What is ABRAXANE used for? Inject the appropriate amount of reconstituted ABRAXANE into an empty, sterile, intravenous infusion bag (plasticized polyvinylidene fluoride). The active agent in ABRAXANE is paclitaxel, a natural product with antitumor activity. The chemical name for paclitaxel is: 10,11-dihydro-5 $\beta$ -[4-oxo-1 $\beta$ -(4-oxo-3-oxo-1 $\beta$ -phenylethoxy)butyl]-7,8,9,10-tetrahydro-6,7,8-trimethoxy-2 $\beta$ -methyl-2 $\beta$ -(propanoic acid)octahydro-1,2-benzoxanthracene-13-one. Paclitaxel is obtained from *Taxus media*. numbness, tingling, pain, or weakness in the hands or feet; nausea. ABRAXANE® for Injectable Suspension (paclitaxel powder for injectable suspension) (nanoparticle, albumin-bound [nab®] paclitaxel) is indicated for: the treatment of metastatic breast cancer. Patient: Dosing volume (mL) = Total dose (mg)/5 (mg/mL). Slowly withdraw the dosing volume of the reconstituted ABRAXANE. ABRAXANE is used to treat the following cancers that have spread to other parts of the body: breast cancer, and pancreatic cancer, when treated. ABRAXANE (nanoparticle albumin-bound paclitaxel) mg powder for injection (suspension) is an albumin nanoparticle form of paclitaxel with a mean particle size of 150 nm. ABRAXANE® for Injectable Suspension (paclitaxel powder for injectable suspension) (nanoparticle, albumin-bound [nab®] paclitaxel) is a cytotoxic anticancer drug and, as such, may cause serious adverse reactions. The most common serious adverse reactions of ABRAXANE in combination with carboplatin for non-small cell lung cancer are anemia (4%) and pneumonia (3%). advanced non-small cell lung cancer. Abbreviated Package Insert. See Product Monograph for complete product information. INDICATIONS. ABRAXANE® for Injectable Suspension (paclitaxel powder for injectable suspension) should be administered under the supervision of a physician experienced in the use of cancer chemotherapeutic agents. ABRAXANE suspension from the vial(s) into a syringe.