



I'm not robot



**I am not robot!**

All items should be timed or counted, except sit to stand and walking through a Objective: To systematically review the evaluation of clinimetric properties and feasibility of the "Modified Parkinson Activity Scale (M-PAS)" and the "Lindop Parkinson's Disease Objectives To assess the validity and inter-rater reliability of the Lindop Parkinson's Disease Mobility Assessment (LPA); a scale developed to gather objective information Sit to Lie Starting position: Sitting on edge of bed/plinth (heightcm), hands on lap, shoes off. Therapist asks patient to lie down. Both M-PAS and LPA are measurement instruments widely used across the world and are specifically designed to evaluate basic functional mobility in patients with PD [21,]. Functional gait Assessment: requiresmetre long/ cm wide walkway Title: Lindop Parkinson's Assessment Scalefor physiotherapists (PDF,KB) Author: User Created Date/25/ PM Guidelines for physiotherapists using Lindop Parkinson's Assessment Scale (Word,KB) Author: user Last modified by: Clare Worrall-Hill Created Date/13/PM Company: sdaht Other titles: Guidelines for physiotherapists using Lindop Parkinson's Assessment Scale (Word,KB) Parkinson's disease is the second most common long-term chronic, progressive, neurodegenerative disease, affecting more thanmillion people worldwide. As soon as patient begins to move, therapist There has been a rising interest in wearable devices for evaluation of movement disorder diseases such as Parkinson's disease due to the limitations in current clinic assessment To assess the validity and inter-rater reliability of the Lindop Parkinson's Disease Mobility Assessment (LPA); a scale developed to gather objective information on gait and bed Title: Lindop Parkinson's Assessment Scalefor physiotherapists (PDF,KB) Author: User Created Date/4/ AM Lindop Parkinson's Physiotherapy Assessment Scale (LPAS) The LPAS is a validated, physiotherapy-specific function measurement tool for use by a physiotherapist to assess the gait and bed mobility of people with Parkinson's. Title: Lindop Parkinson's Assessment Scalefor physiotherapists (PDF,KB) Author: User Created Date/25/ PM Lindop Parkinson's Physiotherapy Assessment Scale (LPAS) Not yet rated The LPAS is a validated, physiotherapy-specific function measurement tool for use by a GUIDELINES FOR PHYSIOTHERAPISTS USING LINDOP PARKINSON'S ASSESSMENT SCALE. M-PAS consists of three domains (Chair Transfer There has been a rising interest in wearable devices for evaluation of movement disorder diseases such as Parkinson's disease due to the limitations in current clinic assessment methods such as Unified Parkinson's Disease Rating Scale Lindop Parkinson's Physiotherapy Assessment Scale (LPAS) Not yet rated The LPAS is a validated, physiotherapy-specific function measurement tool for use by a physiotherapist to assess the gait and bed mobility of people with Parkinson's developed the Modified Parkinson Activity Scale (M-PAS) and Lindop Parkinson's Disease Mobility Assessment (LPA) [11,]. Inability to complete in seconds indicates increased disability and morbidity. The gait section includetasks which are scored out of a total ofwith the higher the score, the better the 5 Times Sit to Stand Test: Assesses likelihood of falling in older adults. Dynamic Gait Index: tests likelihood of falling in older adults. Eight facets of gait tested.