

Schedule An AppointmentoptAmerican Academy of Pediatrics recommends milliliter dosing only. Do not give more thandoses inhours. Acetaminophen Don't give to infants undermonths old unless instructed Recommended dose of MOTRIN® from your healthcare professional: DOSING FOR HEALTHCARE PROFESSIONALS. Ibuprofen (Advil®, Motrin®, or another brand) Give everytohours as needed; always with food. Do not give more thandoses inhours. Weight in pounds (lbs.) Ibuprofen The dosage charts in this handout give the recommended doses for acetaminophen and ibuprofen, medicines that help lower temperature and reduce pain. Pediatric Care Center Acetaminophen and Ibuprofen Dosage Charts Acetaminophen/Ibuprofen Dosage Chart. If you give your child acetaminophen or ibuprofen, be sure to follow these important guidelines. For example, if you take ibuprofen atam you can then take Tylenol atam Dose: Give everytohours as OTC pain relievers dosage chart for adults and childrenyears and older Acetaminophen AMOUNT DOSE & FREQUENCY DAILY LIMIT Acetaminophen regular strength for \* Acetaminophen should be given every hours as needed. Questions? Use only the dosing device (dosing syringe or dosing cup) that came with the product. Do not give more thandoses inhours. \* Do not use with any other medicines that contain acetaminophen Acetaminophen (Tylenol® or another brand) Give every hours as needed. The doses are How to give the right amount of ACETAMINOPHEN (also known as Tylenol) is different depending on which medicine you plan to give. Weight Age LbsMosLbsMosLbsMosLbsYrsLbsYrsLbsYrsLbsYrsLbsYrs. The doses are based on a child's weight. Do not use under agemonths unless advised by your clinician. Do not give more thandoses inhours The dosage charts in this handout give the recommended doses for acetaminophen and ibuprofen, medicines that help lower temperature and reduce pain. \* Do not give more thandoses inhours. Acetaminophen and Ibuprofen Dosing Chart. Do not use any Ibuprofen (Advil®, Motrin®, or another brand) Give everytohours as needed; always with food. Use this chart to determine the proper dose of Always read and follow the label on all TYLENOL® and MOTRIN® products. Acetaminophen. Ibuprofen (Advil®, Motrin®, or another brand) Give everytohours as needed; give with food to lower risk of upset stomach. Ibuprofen Acetaminophen (Tylenol) Dosing Chart. Give everyhours as needed. Important Notes: \* This chart contains two different acetaminophen infant formulations The best way to alternate taking Acetaminophen (example Tylenol) and Ibuprofen (example Advil/Motrin) is to take themhours apart. Give every hours as needed, no more thantimes inhours. inophen (Tylenol® or another. Do not give more thandoses inhours.